

Prevalence of Arthritis Among Adults 18 and Over by State and Region, United States, 2015

	Population with doctor-diagnosed arthritis	Age-standardized prevalence		Population with doctor-diagnosed arthritis	Age-standardized prevalence
Northeast			Midwest		
Maine	332,000	26.4%	Michigan	2,305,000	27.0%
Pennsylvania	2,937,000	25.7%	Missouri	1,372,000	26.8%
Rhode Island	226,000	24.2%	Indiana	1,390,000	25.4%
Vermont	136,000	23.4%	Ohio	2,547,000	25.3%
New Hampshire	282,000	23.0%	Iowa	619,000	23.2%
Massachusetts	1,300,000	22.0%	Kansas	536,000	22.7%
Connecticut	690,000	21.6%	Wisconsin	1,104,000	22.1%
Maryland	1,096,000	21.5%	South Dakota	158,000	21.8%
New York	3,629,000	21.5%	Illinois	2,308,000	21.6%
New Jersey	1,590,000	20.5%	North Dakota	134,000	21.6%
			Nebraska	334,000	21.5%
South			Minnesota	907,000	19.7%
West Virginia	557,000	33.6%	West		
Alabama	1,248,000	30.4%	Oregon	838,000	24.5%
Tennessee	1,630,000	29.4%	Wyoming	116,000	24.1%
Kentucky	1,087,000	29.3%	Montana	216,000	23.9%
Arkansas	672,000	27.1%	Idaho	309,000	23.2%
Mississippi	647,000	26.6%	Washington	1,346,000	22.6%
South Carolina	1,105,000	26.3%	New Mexico	386,000	22.2%
Louisiana	989,000	26.2%	Arizona	1,222,000	21.8%
Oklahoma	813,000	25.7%	Colorado	949,000	21.8%
North Carolina	2,089,000	24.9%	Alaska	117,000	21.5%
Delaware	207,000	24.6%	Utah	407,000	20.8%
Georgia	1,890,000	23.6%	Nevada	477,000	20.1%
Virginia	1,513,000	21.6%	California	5,719,000	18.3%
Florida	4,154,000	21.5%	Hawaii	211,000	17.2%
Dist of Columbia	101,000	19.9%			
Texas	4,055,000	19.0%			
US	54,000,000	23.0%			

States with prevalence above the US rate.

Source: Barbour KE, Moss S, Croft JB, et al. Geographic Variations in Arthritis Prevalence, Health-Related Characteristics, and Management — United States, 2015. MMWR Surveill Summ 2018;67(No. SS-4):1–28. DOI: <http://dx.doi.org/10.15585/mmwr.ss6704a1>